

MATHEMATICS

NOTE ON A DUALITY RELATION OF KAASHOEK

BY

RICHARD K. OLIVER ¹⁾

(Communicated by Prof. A. C. Zaanen at the meeting of January 29, 1966)

The purpose of this note is to give a proof for a correct but incorrectly proven duality relation of KAASHOEK which appears in [2].

First a few preliminary definitions are needed. If T is a linear map with domain $D(T)$ in a vector space X and range $R(T)$ in a vector space Y then we will say that T acts from X to Y ; when it is assumed that $D(T) = X$ we will say simply that T is from X to Y . If T acts from X to Y we define $\alpha(T) = \dim N(T)$, where $N(T)$ is the null space of T , and $\beta(T) = \dim Y/R(T)$, where here and below "dim" denotes the integer k if the vector space in question has finite dimension k and the extended real number $+\infty$ if it is infinite-dimensional. For each pair (T, S) where T and S act from X to Y we define a sequence $(D_n(T, S))_{n \geq 0}$ of subspaces of X as follows: $D_0(T, S) = X$, and for $n > 0$, $D_n(T, S) = S^{-1}TD_{n-1}(T, S)$, where

$$TD_{n-1}(T, S) = \{Tx : x \in D(T) \cap D_{n-1}(T, S)\}$$

and

$$S^{-1}TD_{n-1}(T, S) = \{x : x \in D(S), Sx \in TD_{n-1}(T, S)\},$$

and we set $D(T, S) = \bigcap_{n \geq 0} D_n(T, S)$. If X and Y are normed vector spaces, T acts from X to Y , and $N(T)$ is a closed subspace of X , we define

$$\gamma(T) = \begin{cases} \inf \{\|Tx\|/d(x, N(T)) : x \in D(T), x \notin N(T)\} & \text{if } T \neq 0, \\ +\infty & \text{if } T = 0 \end{cases}$$

where $d(x, N(T))$ denotes the distance from x to $N(T)$.

The following theorem is contained in Theorems 3 and 5 of the paper [1] by KATO:

Theorem 1. *Let X and Y be Banach spaces, T a closed linear map acting from X to Y with $0 < \gamma(T) < \infty$, and S a linear map acting from X to Y with $D(S) \supset D(T)$ and $\|Sx\| \leq \sigma\|x\| + \tau\|Tx\|$ for some non-negative constants σ and τ and all $x \in D(T)$. Then the following statements hold:*

a) *If $N(T) \subset D(T, S)$ then there exists a constant $\varrho > 0$ such that for $|\lambda| < \varrho$, $T - \lambda S$ is closed, $R(T - \lambda S)$ is closed, and $\alpha(T - \lambda S)$ and $\beta(T - \lambda S)$ are constant.*

¹⁾ Written under NASA grant NGR-21-002-054.

b) If $\alpha(T)$ or $\beta(T)$ is finite then there exists a constant $\varrho > 0$ such that for $0 < |\lambda| < \varrho$, $T - \lambda S$ is closed, $R(T - \lambda S)$ is closed, $\alpha(T - \lambda S) = \alpha(T) - k$, and $\beta(T - \lambda S) = \beta(T) - k$ for some non-negative integer k .

On examining Kato's proofs¹⁾ it can be seen that instead of b) the following stronger statement is actually established:

b') If $N(T)/(N(T) \cap D(T, S))$ has finite dimension k then there exists a constant $\varrho > 0$ such that for $0 < |\lambda| < \varrho$, $T - \lambda S$ is closed, $R(T - \lambda S)$ is closed, $\alpha(T - \lambda S) = \alpha(T) - k$, and $\beta(T - \lambda S) = \beta(T) - k$.

In [2] KAASHOEK introduces the quantity $\dim N(T)/(N(T) \cap D(T, S))$ and proves Theorem 1 with statements a) and b') using a method which yields for a) a better value for the constant ϱ than Kato's method; in fact, the value obtained is the best possible.

To prove b') KAASHOEK uses the following duality relation:

$$(1) \quad \dim N(T)/(N(T) \cap D(T, S)) = k \text{ if and only if} \\ \dim N(T^*)/(N(T^*) \cap D(T^*, S^*)) = k,$$

where it is assumed that T and S are continuous linear maps from a Banach space X to a Banach space Y , $R(A)$ is closed, and $k < +\infty$. The proof of this relation is, however, based on a theorem (Theorem 3.3, p. 458 of [2]) which is not true. In this note we present a counterexample to this theorem and then prove a theorem which has (1) as a consequence.

Theorem 3.3 of [2] asserts the following:

Let C , T , and S be continuous linear maps from a Banach space X to a Banach space Y with $N(S) \subset N(C)$, $N(T) \subset D(T, S)$ and $\dim R(C) = l < +\infty$. Then $\dim N(T+C)/(N(T+C) \cap D(T+C, S)) \leq l$.

To see this is not true let $X = Y = l_2$ and define C , T , and S by $Ce_k = 0$, $k \neq 4$, $Ce_4 = -e_2 - e_4$; $Te_k = e_k$, $k \neq 3$, $Te_3 = 0$; and $Se_k = e_{k+1}$, $k \geq 1$; where e_k is the k -th unit vector (δ_{nk}), $n \geq 1$. Let $D = [e_k]_{k \geq 3}$ (= closure of the subspace generated by the vectors e_k , $k \geq 3$) and $R = [e_k]_{k \geq 4}$. It is easily seen that C , T , and S are continuous with $N(S) = [0] \subset N(C)$, $\dim R(C) = 1$, and that $TD = R$, $S^{-1}R = D$, and $[e_3] = N(T) \subset D$, implying $N(T) \subset D(T, S)$. Hence the hypotheses are satisfied with $l = 1$. However, $(T+C)e_k = e_k$, $k \neq 3, 4$, $(T+C)e_3 = 0$, $(T+C)e_4 = -e_2$. Hence $(T+C)X = [e_k]_{k \neq 3, 4}$,

$$S^{-1}(T+C)X = [e_k]_{k \neq 2, 3},$$

and $N(T+C) = [e_2 + e_4, e_3]$. Thus $N(T+C) \cap S^{-1}(T+C)X = [0]$, implying $\dim N(T+C)/(N(T+C) \cap D(T+C, S)) = 2 > l = 1$.

The relation (1) is a consequence of the following theorem:

¹⁾ Specifically it can be seen that Kato's Theorem 4, p. 307 of [1], holds if the hypothesis " $\alpha(A)$ or $\beta(A)$ is finite" is replaced by " $\dim N(A)/(N(A) \cap D(A, B)) = k < +\infty$ " and that then the " r " of this theorem is equal to k ; with this observation, statement b') follows by the proof of Theorem 5, p. 315 of [1].

Theorem 2. *Let T and S be continuous linear maps from a Banach space X to a Banach space Y with $R(T)$ closed. Then for all $n \geq 0$,*

$$\dim N(T)/(N(T) \cap D_n(T, S)) = \dim N(T^*)/(N(T^*) \cap D_n(T^*, S^*)).$$

Proof. Set $D_n = D_n(T, S)$, $D_n^* = D_n(T^*, S^*)$, and define a sequence $(N_n)_{n \geq 0}$ of subspaces of X as follows: $N_0 = \{0\}$, and for $n > 0$, $N_n = T^{-1}SN_{n-1}$. We first show

$$(2) \quad \dim N_n/(N_n \cap D_1) = \dim N_1/(N_1 \cap D_n), \quad n \geq 0.$$

Since

$$\begin{aligned} \dim N_n/(N_n \cap D_1) &= \dim (N_n + D_1)/D_1 = \dim (N_n + D_1)/(N_{n-1} + D_1) + \\ &\quad \dots + \dim (N_1 + D_1)/(N_0 + D_1), \end{aligned}$$

and

$$\begin{aligned} \dim N_1/(N_1 \cap D_n) &= \dim (N_1 \cap D_0)/(N_1 \cap D_1) + \dots \\ &\quad + \dim (N_1 \cap D_{n-1})/(N_1 \cap D_n), \end{aligned}$$

it is enough to show

$$\begin{aligned} \dim (N_{k+1} + D_1)/(N_k + D_1) &= \dim (N_1 \cap D_k)/(N_1 \cap D_{k+1}), \\ 0 \leq k \leq n-1. \end{aligned}$$

Let $(x_i^0)_{1 \leq i \leq n}$ be a sequence in $N_1 \cap D_k$ linearly independent modulo $N_1 \cap D_{k+1}$. Since it is in D_k there exist sequences $(x_i^j)_{1 \leq i \leq n} \subset D_{k-j}$, $1 \leq j \leq k$, such that

$$(3) \quad Sx_i^0 = Tx_i^1, Sx_i^1 = Tx_i^2, \dots, Sx_i^{k-1} = Tx_i^k.$$

Since it is also in N_1 we have from (3) that $(x_i^k)_{1 \leq i \leq n}$ is in N_{k+1} . Put $x^j = \sum_{i=1}^n \alpha_i x_i^j$, $0 \leq j \leq k$, and suppose $x^k \in N_k + D_1$. Then $x^k = n^k + d^1$ for some $n^k \in N_k$ and $d^1 \in D_1$, and hence $Sx^{k-1} = Tx^k = Tn^k + Td^1 = Sn^{k-1} + Td^1$ for some $n^{k-1} \in N_{k-1}$. Thus $S(x^{k-1} - n^{k-1}) = Td^1$, i.e., $x^{k-1} = n^{k-1} + d^2$ for some $d^2 \in D_2$. Proceeding in this way we get $x^1 = n^1 + d^k$ for some $n^1 \in N_1$ and $d^k \in D_k$ and hence $Sx^0 = Tx^1 = Td^k$, i.e., $x^0 \in D_{k+1}$. Since $(x_i^0)_{1 \leq i \leq n}$ is linearly independent modulo $N_1 \cap D_{k+1}$ it follows that $\alpha_1 = \dots = \alpha_n = 0$, proving $(x_i^k)_{1 \leq i \leq n}$ is linearly independent modulo $N_k + D_1$. Thus

$$\dim (N_{k+1} + D_1)/(N_k + D_1) \geq \dim (N_1 \cap D_k)/(N_1 \cap D_{k+1}).$$

To get the opposite inequality let $(x_i^0)_{1 \leq i \leq n}$ be a sequence in N_{k+1} linearly independent modulo $N_k + D_1$. Then there exist sequences

$$(x_i^j)_{1 \leq i \leq n} \subset N_{k-j+1}, \quad 1 \leq j \leq k$$

such that

$$(4) \quad Tx_i^0 = Sx_i^1, Tx_i^1 = Sx_i^2, \dots, Tx_i^{k-1} = Sx_i^k.$$

From (4) it follows that $(x_i^k)_{1 \leq i \leq n}$ is in D_k . Put $x^j = \sum_{i=1}^n \alpha_i x_i^j$, $0 \leq j \leq k$,

and suppose $x^k \in D_{k+1}$. Then $Tx^{k-1} = Sx^k = Td^k$ for some $d^k \in D_k$, implying $x^{k-1} - d^k = n^1$ for some $n^1 \in N_1$. Similarly,

$$Tx^{k-2} = Sx^{k-1} = Sn^1 + Sd^k = Sn^1 + Td^{k-1}$$

for some $d^{k-1} \in D_{k-1}$ and hence $x^{k-2} \in D_{k-1} + N_2$. Repeating this procedure we finally get $x^0 \in N_k + D_1$, implying $\alpha_1 = \dots = \alpha_n = 0$. Thus

$$\dim (N_{k+1} + D_1) / (N_k + D_1) \leq \dim (N_1 \cap D_k) / (N_1 \cap D_{k+1}).$$

We now show $D_n^* \subset (SN_n)^\perp$, $n \geq 0$. This is obvious for $n=0$. Suppose $f \in D_{n+1}^*$, $x \in N_{n+1}$, $g \in D_n^*$ is such that $S^*f = T^*g$, and $y \in N_n$ is such that $Tx = Sy$. Then $f(Sx) = (S^*f)x = (T^*g)x = g(Tx) = g(Sy)$. Thus $D_n^* \subset (SN_n)^\perp$ implies $D_{n+1}^* \subset (SN_{n+1})^\perp$, and the relation $D_n^* \subset (SN_n)^\perp$, $n \geq 0$, follows by induction.

With the relations $D_1 = S^{-1}R(T)$ and $R(T)^\perp = N(T^*)$, and the hypothesis that $R(T)$ is closed it is not difficult to show that

$$(5) \quad \dim N_n / (N_n \cap D_1) = \dim (R(T) + SN_n) / R(T) = \\ = \dim N(T^*) / (N(T^*) \cap (SN_n)^\perp).$$

If for some n (and hence for all larger n in view of $N_{n+1} \supset N_n$)

$$\dim N(T^*) / (N(T^*) \cap (SN_n)^\perp) = +\infty$$

let p be the least such n ; otherwise set $p = +\infty$. Since, as we have already shown, $D_n^* \subset (SN_n)^\perp$ for all n ,

$$(6) \quad \dim N(T^*) / (N(T^*) \cap (SN_n)^\perp) = \dim N(T^*) / (N(T^*) \cap D_n^*)$$

holds for $n \geq p$. For $n < p$ we will prove (6) by showing $(SN_n)^\perp \subset D_n^*$, and thus complete the proof in view of (2) and (5). This is clear for $n=0$. Assume $(SN_n)^\perp \subset D_n^*$ and let $f \in (SN_{n+1})^\perp$. If there is a $g \in (SN_n)^\perp$ such that $T^*g = S^*f$, then from $(SN_n)^\perp \subset D_n^*$ and the definition of D_{n+1}^* it follows that $f \in D_{n+1}^*$, i.e., $(SN_{n+1})^\perp \subset D_{n+1}^*$. We now find such a g . Since $N(T) = N_1 \subset N_{n+1}$ we have $S^*f(N(T)) \subset S^*f(N_{n+1}) = \{0\}$. We can therefore define a linear functional g' on the range $R(T)$ by the relation $g'(Tx) = (S^*f)x$; g' is continuous because

$$|g'(Tx)| = \inf \{ |g'(T(x+x'))| : x' \in N(T) \} \\ \leq \inf \{ \|S^*f\| \cdot \|x+x'\| : x' \in N(T) \} \leq \|S^*f\| \cdot \gamma(T)^{-1} \cdot \|Tx\|.$$

Since $\dim (R(T) + SN_n) / R(T)$ is finite, there is a finite-dimensional subspace M of Y such that $SN_n = M \oplus (SN_n \cap R(T))$. Let P be the projection in the Banach space $M \oplus R(T)$ with null space M and range $R(T)$ and define $g'' = g' \circ P$. If $x \in N_n$ and $Sx \in R(T)$ we have $Sx = Tx'$ for some $x' \in N_{n+1}$, so that $g''(Sx) = g''(Tx') = g'(Tx') = f(Sx') = 0$. Since g'' annihilates

M we thus have $g''(SN_n) = \{0\}$. Finally, since g'' is continuous,

$$(\|g''\| \leq \|g'\| \cdot \|P\|),$$

there is a $g \in Y^*$ extending g'' , and any such g obviously has the required properties. This completes the proof.

I would like to thank Dr. KAASHOEK for pointing out some obscurities in the original version of the proof of Theorem 2.

University of Maryland

REFERENCES

1. KATO, T., Perturbation theory for nullity, deficiency and other quantities of linear operators, *J. d'Analyse Math.* **6**, 261–322 (1958).
2. KAASHOEK, M. A., Stability theorems for closed linear operators, *Proc. Kon. Ned. Akad. v. Wetensch.*, Amsterdam, **A 68**, 452–466 (1965).